

Varicella (chickenpox)

CLINICAL CASE DEFINITION

An illness with acute onset of diffuse (generalized) maculo-papulo-vesicular rash without other apparent cause.

CASE CLASSIFICATION

- Probable:** a case that meets the clinical case definition, is not laboratory confirmed, and is not epidemiologically linked to another probable or confirmed case.
- Confirmed:** a case that is laboratory confirmed or that meets the clinical case definition and is epidemiologically linked to a confirmed or probable case.

Acceptable laboratory confirmation is any of the following:

- Isolation of varicella virus from a clinical specimen, OR
- Varicella antigen detected by direct fluorescent antibody (DFA) test, OR
- Varicella-specific nucleic acid detected by polymerase chain reaction (PCR), OR
- Significant rise in serum anti-varicella immunoglobulin G (IgG) antibody level by any standard serologic assay.

Comments:

- Two probable cases that are epidemiologically linked would be considered confirmed, even in the absence of laboratory confirmation.
- Occasionally, a vaccinated person does not get full immunity and may still get the disease. This is known as “breakthrough disease.” Varicella breakthrough disease cases are usually mild, with fewer than 50 skin lesions and a shorter duration of illness. The rash may also be atypical in appearance (maculopapular with few or no vesicles).
- Laboratory confirmation is now encouraged. Although laboratory confirmation of cases in the past was rarely sought, fewer clinicians now have direct diagnostic experience since varicella incidence has substantially declined as a result of vaccination. See LABORATORY SPECIMENS: PROCEDURES AND CONSIDERATIONS, below for further information.

TRANSMISSION

Person-to-person via direct contact or droplet or airborne spread of respiratory tract secretions or vesicle fluid of patients; also from vesicle fluid of persons with herpes zoster (shingles). Chickenpox is highly communicable.

INCUBATION PERIOD

14 – 16 days, range 10 – 21 days. See [Varicella Timeline](#), below.

PERIOD OF COMMUNICABILITY

From 1-2 days before onset of rash until all lesions have crusted (for breakthrough cases who may develop lesions that don't crust: until lesions are fading or until no new lesions occur, whichever is later).

REPORTING/INVESTIGATION

Health care providers, schools, day care providers and camps should report cases/suspect cases of varicella to local health department serving the residence of the case.

Local health department role/responsibilities:

- ◆ Contact case/guardian and health care provider.
- ◆ Determine if case meets clinical case definition.
- ◆ If definition met (probable or confirmed cases), or otherwise suspected as a case, investigate using CDC surveillance worksheet and control guidelines below.
- ◆ Assist with coordination of specimen collection and coordination if public health lab resources (MDCH, CDC, etc) are used
- ◆ Report/ensure reporting of case to the Michigan Disease Surveillance System (MDSS). [CDC Varicella Surveillance Worksheet](#) may be helpful in field investigation to collect and capture data. At a minimum, obtain basic demographic information, immunization history (number of doses and dates) from provider record or MI Care Improvement Registry (MCIR - state immunization registry), and an estimate of the number of lesions, which serves as a proxy for disease severity. Number of lesions can be approximated as follows:
 - **Less than 50 lesions** –the lesions can be easily counted within 30 seconds
 - **50-249 lesions** – the person's hand can be placed between the lesions without touching a lesion
 - **250-499 lesions** – the person's hand cannot be placed between the lesions without touching a lesion
 - **More than 500 lesions** – in this case, the lesions are clumped so closely together that it is difficult to see normal skin.
- ◆ Update the MDSS record in a timely manner with new or additional info as it becomes available. Finalize MDSS record when case investigation is complete.
- ◆ In the event of death, obtain and send copies of hospital discharge summary, death certificate, and autopsy report to MDCH Immunization Division.
- ◆ Outbreak reporting: Varicella outbreaks consisting of 5 or more cases in a group activity setting (school, daycare, camp, etc) should be reported to MDCH VPD Surveillance Coordinator at 517-335-8159; reporting of outbreaks consisting of fewer cases is also encouraged . Complete the [varicella outbreak reporting worksheet](#) (below) and submit to MDCH when outbreak has concluded. Information needed is total number of cases and number within certain age-groups, dates of first case onset and last case onset, setting (school, day care, etc), number previously vaccinated (and number of doses, i.e. number with 1 dose, number with 2 doses) by certain age-groups, number never vaccinated, median age and age range, and # confirmed by laboratory testing

LABORATORY CONFIRMATION

Lab confirmation of varicella cases is encouraged now as overall incidence has greatly declined and clinical experience with the disease becomes more rare. Confirmation is strongly recommended to confirm the diagnosis in severe or unusual cases and for fatal cases. **Laboratory confirmation for varicella is defined as:**

- Isolation of varicella virus from a clinical specimen, or
- Detection of varicella antigen or nucleic acid by direct fluorescent antibody (DFA) test, or
- Varicella-specific nucleic acid detected by polymerase chain reaction (PCR) test, or
- Significant rise in serum varicella immunoglobulin G (IgG) antibody level by any standard serologic assay
- Note: varicella IgM antibody testing at commercial laboratories is not recommended for confirmation because the methods for these tests lack sensitivity and specificity; however, a positive varicella IgM test **performed by CDC's direct capture IgM assay** will be considered confirmatory.

Varicella lab tests are available commercially. See additional information under [LABORATORY SPECIMENS: PROCEDURES AND CONSIDERATIONS](#), below

IMMUNITY/SUSCEPTIBILITY

Individuals should be considered immune (protected against) varicella if they meet one or more of the following conditions:

1. Documentation of age-appropriate vaccination:
 - a. Preschool-aged children ≥ 12 months of age: one (1) dose¹
 - b. School-aged children, adolescents, and adults: two (2) doses¹
2. Laboratory evidence of immunity² or laboratory confirmation of disease
3. Born in the US before 1980³
4. A healthcare provider diagnosis of varicella or healthcare provider verification of history of varicella disease⁴
5. History of herpes zoster based on healthcare provider diagnosis.

¹ The minimum interval between doses is 28 days for persons ≥ 13 years of age and is 3 months for children 12 months through 12 years of age; however, for children who have received their first dose before age 13 years and the interval between the two doses was at least 28 days, the second dose is considered valid and need not be repeated.

² Commercial assays can be used to assess disease-induced immunity, but they lack adequate sensitivity to detect reliably vaccine-induced immunity (may yield false negative results).

³ For healthcare providers and pregnant women, birth before 1980 should not be considered evidence of immunity.

⁴ Verification of history or diagnosis of typical disease can be done by any healthcare provider (e.g., school or occupational clinic nurse, nurse practitioner, physician assistant, physician). For people reporting a history of or presenting with atypical and/or mild cases, assessment by a physician or their designee is recommended and one of the following should be sought: a) an epidemiologic link to a typical varicella case or b) evidence of laboratory confirmation, if laboratory testing was performed at the time of acute disease. When such documentation is lacking, people should not be considered as having a valid history of disease, because other diseases may mimic mild atypical varicella.

CONTROL MEASURES

- ◆ Exclude cases or suspected cases from group activity settings (e.g. schools, day-care centers, work places, camps) until all lesions have crusted. Instruct cases/suspect cases to avoid exposing other persons.
Note: Vaccinated persons with varicella (i.e. “breakthrough” cases) may develop lesions that do not crust (that is, macules and papules only, no vesicles). These

persons are considered no longer contagious once lesions have faded (skin lesions in the process of resolving), or once no new lesions occur, whichever is later.

- ◆ Identify exposed contacts and determine if they are immune or susceptible.
- ◆ Exposed, susceptible persons should be vaccinated as soon as possible; post-exposure vaccination with varicella vaccine given within 3 days of exposure may prevent illness or modify severity of disease (studies indicate 70% - 100% effectiveness). Post-exposure prophylaxis use of vaccine up to 5 days after exposure may also be effective.
- ◆ Varicella immune globulin (VZIG/VariZIG™) for postexposure prophylaxis of varicella should be considered in persons at high risk for severe disease who lack evidence of immunity to varicella and are ineligible for varicella vaccine:
 - VariZIG™ (Cangene Corporation, Winnipeg, Canada) is the only varicella zoster immune globulin preparation available in the United States
 - VariZIG™ is available in the United States through an investigational new drug (IND) application expanded access protocol
 - VariZIG™ can be obtained by health-care providers from the sole-authorized U.S. distributor, FFF Enterprises (Temecula, California), by calling 800-843-7477 at any time or by contacting the distributor online at <http://www.fffenterprises.com>
 - VariZIG™ should be administered intramuscularly as directed by the manufacturer and given as soon as possible, ideally within 4 days of exposure, but may be used up to 10 days after
 - The recommended dose is 125 IU/10 kg of body weight, up to a maximum of 625 IU (five vials) (VariZIG is supplied in 125-IU vials)
 - Patient groups recommended by ACIP to receive VariZIG™ include the following:
 - Immunocompromised patients without evidence of immunity.
 - Neonates whose mothers have signs and symptoms of varicella around the time of delivery (i.e., 5 days before to 2 days after).
 - Hospitalized premature infants born at ≥28 weeks of gestation whose mothers do not have evidence of immunity to varicella.
 - Hospitalized premature infants born at <28 weeks of gestation or who weigh ≤1,000 g at birth, regardless of their mothers' evidence of immunity to varicella.
 - Pregnant women without evidence of immunity.
 - Additional information on VariZIG™ is available at http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6228a4.htm?s_cid=mm6228a4_w
- ◆ Outbreaks in group-activity settings (e.g. schools, day-care centers, work place, camps): Exposed persons who cannot provide documentation of varicella immunity should be excluded until 21 days after the last identified case; this measure is advisable when 2 or more varicella (chickenpox) cases have occurred, but it may also be considered in the instance of a single case.
 - In general, an excluded person may be re-admitted to the activity setting/institution immediately upon getting vaccinated or providing other acceptable documentation of immunity; however, such a re-admittance policy may be modified depending upon the circumstances involved.
 - If the vaccine dose given was the first varicella vaccine dose, the person may *conditionally* return to school, group program, etc., but the second dose of vaccine should be scheduled using the age-appropriate minimal interval (3

months for persons 1 – 12 years of age, 28 days for persons 13 years and older).

- *Outbreaks in settings with children between 12 months and 4 years of age:* While routinely the 2nd dose of varicella vaccine is not given until 4 – 6 years of age, in outbreak situations involving day care, pre-school, and other settings with children under 4 years of age consideration should be given to requiring the 2nd dose as a control measure, using appropriate minimum intervals between doses.
- ◆ Enhance surveillance in the affected setting and community.
- ◆ Guidance for outbreak investigation and response is available from MDCH; additional guidance is available in the [CDC varicella outbreak manual \(http://www.cdc.gov/chickenpox/outbreaks/control-investigation.html\)](http://www.cdc.gov/chickenpox/outbreaks/control-investigation.html)
- ◆ Management of persons with herpes zoster (shingles) in school or group program:
 - Immunocompetent persons with shingles can remain at school as long as the lesions can be completely covered. Persons with shingles should be careful about personal hygiene, wash their hands after touching their lesions and also avoid close contact with others. If the lesions cannot be completely covered and close contact avoided, children and staff should be excluded from the school setting until lesions have crusted over. If a person has disseminated shingles, he or she should be excluded from school until lesions have crusted over (similar to the management of varicella case-patients).

LABORATORY PROCEDURES AND CONSIDERATIONS

Laboratory confirmation of varicella is increasingly important as incidence continues to decline and health care providers have less familiarity and clinical diagnostic experience with the disease. Moreover, breakthrough disease often involves an atypical rash and presentation (fewer lesions, vesicles absent, and milder illness than typical varicella cases). Laboratory confirmation is also encouraged for severe cases (e.g. involving hospitalization or death). Confirmation of at least one case, and preferably 3-5 cases, is recommended in outbreaks.

PCR (polymerase chain reaction) tests are the method of choice for rapid clinical diagnosis and confirmation. Specimens for VZV PCR include swabs of fluid from unroofed vesicles and/or crusts.

DFA tests are an alternate method of confirmation; these are not as sensitive as PCR. Appropriate specimens for DFA are a scraping or swab from the base of open vesicles.

Varicella diagnostic testing is not available through MDCH; consult with commercial laboratories.



MI Varicella Outbreak Reporting Worksheet


For varicella outbreaks of 5 or more cases, please complete this worksheet and send to MDCH. Outbreaks with fewer cases are also encouraged to be reported.

Date of report to MDCH __/__/__

County		Person reporting		Phone		Email		Fax						
Outbreak Name	Outbreak Dates (onset first case; onset last case)	Setting (eg. elementary school, day care, camp)	Total Number of cases	Number of cases in each age group		Number of cases in each lesion category		Vaccination status Of cases				Vaccination coverage in setting		# lab-confirmed cases in outbreak
								#Vax		#Un-vax		1 dose %	2 dose %	
								1 dose	2 dose	0 doses	Prior disease / immune			
				<1	15-19	<50								
				1-4	≥20	50-249								
				5-9	Unk	250-499								
				10-14		≥500								

Send form to
MDCH, Division of Immunization
Attn: VPD Surveillance Coordinator
PO Box 30195
Lansing, MI 48909
Fax 517-335-9855

Varicella* timeline diagram



* This diagram applies only to chickenpox. Shingles is also caused by varicella zoster virus, but is much less infectious.
† Cases may be asymptomatic or very mild, especially in vaccinated individuals. Such cases can still transmit disease.

Sources: APHA Control of Communicable Diseases Manual, AAP Red Book, CDC Pink Book, CDC VPD surveillance manual